



Case Report ●●●

Ossification of vascular pedicle in fibula free flap: a case report

Ossificação do pedículo vascular de um retalho livre de fíbula: relato de caso

MARIA CECÍLIA CLOSS ONO¹

ANNE KAROLINE GROTH²

JOÃO MANOEL MOREIRA³

ALFREDO BENJAMIN DUARTE DA SILVA⁴

IVAN MALUF JUNIOR⁵

■ ABSTRACT

Introduction: The use of fibula flaps for the reconstruction of craniomaxillofacial defects has many advantages, including the low morbidity of the donor area, good bone quality for use of osseointegrated implants, and the possibility to include a skin island, when indicated. During the dissection of the flap, a muscle "cuff" and a periosteal strip are usually included near the region of the vascular pedicle. The osteogenic potential of the transplanted periosteum has been the object of studies. **Case report:** A 15-year-old male patient underwent microsurgical reconstruction using a fibula flap for a mandibular defect caused by the resection of a bone sarcoma. He developed increased volume and bone consistency in the cervical region next to the area where a cervicotomy was performed for vascular anastomosis. Imaging examinations showed the characteristics of the bone mass. He then underwent a new cervicotomy and mass exploratory surgery because bone tissue formation was observed at the site of vascular anastomosis. Anatomopathological examination of the specimen showed bone tissue formation next to the periosteal flap. **Discussion:** During fibula flap dissection, osteotomy is performed a few centimeters from the knee joint to facilitate the dissection of the vascular pedicle in the region of the popliteal fossa. Then, the vascular pedicle is surrounded by a muscle cuff and periosteal strip. This maintains its osteogenic capacity, which can be activated according to the stimulus of the area. Although periosteal ossification of the vascular pedicle in fibula free flaps is a rare event, it has been reported in different centers.

Keywords: osteogenic capacity; fibula free flaps; ossification.

■ RESUMO

Introdução: Várias são as vantagens da utilização de retalhos fibulares para as reconstruções de defeitos craniomaxilofaciais, incluindo a baixa morbidade da área doadora, boa qualidade óssea possibilitando a realização de implantes osteointegrados quando indicados, além da possibilidade de inclusão de uma ilha de pele quando indicado. Durante a dissecação do retalho, próximo à região do pedículo vascular, normalmente inclui-se um cuff muscular e uma faixa de periósteo. O potencial

Institution: This study was performed at the Erasto Gaertner Hospital, Parana League against Cancer, Curitiba, PR.

Article submitted: December 2, 2012.

Article accepted: March 10, 2013.

DOI: 10.5935/2177-1235.2014RBCP0037

1 – Plastic and craniomaxillofacial surgeon, specialist member of the Brazilian Society of Plastic Surgery and Brazilian Association of Craniomaxillofacial Surgery, Plastic Surgery Service of Erasto Gaertner Hospital, Curitiba, PR, Brazil; – associate professor at the Plastic and Reconstructive Surgery Service of the Hospital das Clínicas of UFPR.

2 – Plastic surgeon, member of the Brazilian Society of Plastic Surgery, Plastic Surgery Service of Erasto Gaertner Hospital, Curitiba, PR, Brazil; – Plastic Surgery Service of Erasto Gaertner Hospital.

3 – Plastic surgeon, specialist member of the Brazilian Society of Plastic Surgery, specialist in Oncology Reconstructive Surgery at Erasto Gaertner Hospital, Curitiba, PR, Brazil; – specialist member of the Brazilian Society of Plastic Surgery.

4 – Plastic surgeon, member of the Brazilian Society of Plastic Surgery – chief of the Plastic Surgery Service of Erasto Gaertner Hospital, Curitiba, PR, Brazil.

5 – General surgeon and plastic surgery resident at the Hospital das Clínicas of UFPR.

osteogênico do perióstio transplantado tem sido objeto de estudo. **Relato de caso:** paciente de 15 anos, submetido à reconstrução microcirúrgica com um retalho fibular para um defeito mandibular pós-resssecção de um sarcoma ósseo. Evoluiu com aumento de volume, de consistência óssea na região cervical próximo à cervicotomia realizada para anastomose vascular. Exames de imagem mostravam características ósseas da massa. Foi então submetido à nova cervicotomia e exploração da massa, sendo observada uma formação de tecido ósseo no local da anastomose vascular. Exame anatomopatológico da peça mostrava formação de tecido ósseo adjacente ao retalho periostal. **Discussão:** Durante a dissecação do retalho fibular, a osteotomia é realizada a alguns centímetros da articulação do joelho, isto a fim de facilitar a dissecação do pedículo vascular na região do oco poplíteo. O pedículo vascular fica então envolto por uma cuff muscular e por uma tira de perióstio. Este mantém sua capacidade osteogênica, que pode ser ativada de acordo com o estímulo do local. A ossificação do perióstio do pedículo vascular de retalhos livres de fíbula permanece um evento raro, porém relatado por centros diferentes.

Descritores: Capacidade osteogênica; Retalho livre de fíbula; ossificação.

INTRODUCTION

The use of fibula free flaps is a procedure of choice for the reconstruction of large mandibular defects, especially those involving the anterior mandibular region and occurring after a large tumor resection. Its bone length and rich vascularization allow jaw remodeling by using multiple osteotomies^{1,2}.

The use of fibula flaps for the reconstruction of craniomaxillofacial defects has many advantages, including the low morbidity of the donor area, good bone quality for use of osseointegrated implants, and the possibility to include a skin island, when indicated, in addition to the possibility of having two surgical teams working simultaneously³.

During the dissection of the flap, a muscle "cuff" and a periosteal strip are usually included next to the region of the vascular pedicle. The osteogenic potential of the transplanted periosteum has been the object of several studies^{3,4}; however, its clinical observation in fibula osteomyocutaneous flaps has been described as a rare event⁵.

The objective of this study is to relate a case of periosteal ossification of the vascular pedicle in a fibula flap and present a brief literature review.

CASE REPORT

A 15-year-old male patient underwent microsurgical reconstruction with a fibula flap for a mandibular defect caused by resection of a bone sarcoma.

Nine months after surgery, the patient developed increased volume and bone consistency in the cervical region next to the area where a cervicotomy was performed for vascular anastomosis. Imaging examinations showed the characteristics of the bone mass. Then, he underwent a new cervicotomy and mass exploratory surgery because bone tissue formation was observed at the site of vascular anastomosis.

Anatomopathological examination of the specimen showed bone tissue formation next to the periosteal flap.

DISCUSSION

Although periosteal ossification of the vascular pedicle in fibula free flaps is a rare event, it has been reported to occur in different centers. It was initially reported in 2003⁶, and four more cases were reported in 2008⁷.

During fibula flap dissection, osteotomy is performed a few centimeters from the knee joint to facilitate the dissection of the vascular pedicle in the region of the popliteal fossa. This osteotomy is performed after periosteal detachment, to preserve the longitudinal vascularization along the flap (vascularization through the periosteum)⁸. Then, the vascular pedicle is surrounded by a muscle cuff and periosteal strip. This maintains its osteogenic capacity³, which can be activated according to the stimulus of the area.

The ability of the periosteum to regenerate a new bone can be observed especially in younger patients and in revascularized flaps. This characteristic, combined with direct contact with the bone, allows the formation of a new bone along the pedicle. Therefore, it becomes necessary to modify the surgical technique while reducing excess fibula bone, removing the periosteum with the bone to avoid the complication described above⁷.

In cases in which the primary disease is a tumor, the diagnosis can be confused with recurrence. Depending on the area of the pedicle and ossification, the patient may present symptoms or be completely asymptomatic⁹.

The osteogenic potential of the periosteum exists with fibula free flap transfer and, depending on the location, it may have significant consequences such as intense pain, increased local volume, and even trismus⁶. Ossification of the vascular pedicle is unusual and probably underdiagnosed. The

complications are rare, and the surgical removal of ossification should be reserved for patients with symptoms¹⁰.

REFERENCES

1. Braga-Silva J, Martins PD, Román JA, Gehlen D. Reconstrução do segmento ósseo mandibular: comportamento dos implantes ósseo-integrados nos retalhos vascularizados de crista ilíaca e fibula; *Rev Soc Bras Cir Plást.* 2005;20(3):176-81.
2. Cardoso IF, Sbalchiero JC, Batista AS, Ohana BM, Chedid R, Cardoso GF, et al. Use of fibular osteocutaneous flap in microsurgical reconstruction of complex mandibular defects. *Rev Bras Cir Plást.* 2011;26(1):42-7.
3. Asamura S, Ikada Y, Matsunaga K, Wada M, Isogai N. Treatment of orbital floor fracture using a periosteum-polymer complex. *J Craniomaxillofac Surg.* 2010;38(3):197-203.
4. Sakata Y, Ueno T, Kagawa T, Kanou M, Fujii T, Yamachika E, et al. Osteogenic potential of cultured human periosteum-derived cells – a pilot study of human cell transplantation into a rat calvarial defect model. *J Craniomaxillofac Surg.* 2006;34(8):461-5.
5. González-García R, Manzano D, Ruiz-Laza L, Moreno-García C, Monje F. The rare phenomenon of vascular pedicle ossification of free fibular flap in mandibular reconstruction. *J Craniomaxillofac Surg.* 2011;39(2):114-8.
6. Smith RB, Funk GF. Severe trismus secondary to periosteal osteogenesis after fibula free flap maxillary reconstruction. *Head Neck.* 2003;25(5):406-11.
7. Autelitano L, Colletti G, Bazzacchi R, Biglioli F. Ossification of vascular pedicle in fibular free flaps: a report of four cases. *Int J Oral Maxillofac Surg.* 2008;37(7):669-71.
8. Urken ML, Buchbinder D, Costantino PD, Sinha U, Okay D, Lawson W, et al. Oromandibular reconstruction using microvascular composite flaps: report of 210 cases. *Arch Otolaryngol Head Neck Surg.* 1998;124(1):46-55.
9. Acartürk TO, Aslaner EE. Periosteal ossification from the vascular pedicle of a free fibular flap. *J Craniofac Surg.* 2011;22(6):e29-32.
10. Myon L, Ferri J, Genty M, Raoul G. Consequences of bony free flap's pedicle calcification after jaw reconstruction. *J Craniofac Surg.* 2012;23(3):872-7.

Corresponding Author:

Maria Cecília Closs Ono

Rua Paulo Martins, 158, cs 03, CEP 80710010

Fone/Fax +55 041 30763418

E-mail: mccono@gmail.com